

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/296701534>

Presacral Schwannoma Resected Via Anterior Trans Peritoneal Approach: A Case Report

Article · March 2016

CITATIONS

0

READS

25

3 authors, including:



Vishwas Pai

Pai Onco Care Centre, Hubli,, Karnataka, India.

58 PUBLICATIONS 112 CITATIONS

SEE PROFILE

Presacral Schwannoma Resected Via Anterior Trans Peritoneal Approach: A Case Report

Case Report

Abstract

Presacral tumors are rare. Neurogenic tumors in this area are even rarer with only case report and short case series being reported in literature. Most of these tumors remain asymptomatic till they reach size large enough to cause pressure symptoms. Magnetic resonance imaging (MRI) of the pelvis is the most preferred imaging modality for preoperative diagnosis as well treatment planning. Complete surgical resection is the treatment of choice. Approach to surgical resection depends on the type of the tumour. After complete excision of the tumor, recurrence is rare. Up to 10-50% recurrence has been reported in literature after incomplete resections. Adjuvant treatment is not recommended even in incomplete resections or recurrences. We are presenting a case of type 3 presacral schwannoma resected by open, anterior, trans peritoneal approach.

Keywords: Presacral schwannoma; Neurogenic tumors; Anterior transperitoneal approach

Volume 3 Issue 2- 2015

Vishwas D Pai^{1*}, Ashutosh Pawale² and Suvarna Ravindranath³

¹Department of Surgical Oncology, Kerudi Cancer Hospital and Research Centre, Bagalkot, Karnataka, India

²Department of Radiodiagnosis, Kerudi Cancer Hospital and Research Centre, Bagalkot, Karnataka, India

³Department of Pathology, SRL Diagnostics, Bangalore, Karnataka, India

***Corresponding author:** Vishwas D Pai, Department of Surgical Oncology, Kerudi Cancer Hospital and Research Centre, Bagalkot 587101, Karnataka, India, Tel: +91-9449333502; E-mail: vishpai88@gmail.com

Received: October 28, 2015 | **Published:** December 29, 2015

Abbreviations: CECT: Contrast Enhanced Computed Tomography; MRI: Magnetic Resonance Imaging; CT: Computed Tomography

Introduction

Schwannomas are the most common types of peripheral nerve neoplasms arising from the Schwann cells of the neural sheath. Although they may occur in any part of the body, only 6% are found in the presacral area [1]. Exact incidence is difficult to estimate since majority of these remain asymptomatic. Because of the location, these tumors can grow to very large size before being diagnosed. These tumors offer significant surgical challenge because of the narrow working space available and proximity to the vital structures including external iliac vessels as well as the ureter. In addition, abundant vascularity of the presacral plexus of vessels can lead to significant blood loss. Accurate delineation of the type of the tumor on preoperative imaging studies is essential to determine the appropriate approach for these rare tumors. We are presenting a case of presacral schwannoma resected via an open, intraperitoneal approach.

Case Report

A 40 year old gentleman presented with pain in the lower abdomen of 1 year duration. On evaluation with contrast enhanced computed tomography of abdomen (CECT), a soft tissue mass of size 10 x 8 x 6 cm was detected in the presacral area. A Magnetic resonance imaging (MRI) of the pelvis confirmed it as dumb-bell shaped tumor with no intra spinal extension representing a nerve sheath tumour (Figure 1a). Intra operatively, right ureter was mobilized first and the tumor was dissected from right external iliac artery and vein. Tumor was then separated from the sacrum by sharp and blunt dissection. Excision of the tumor was completed after clipping the 5th lumbar nerve root on the right side (Figure 1b). Duration of the surgery was 100

minutes and there was 100 ml of blood loss. Post operative course was uneventful and he was discharged on 4th post operative day. Histopathological examination of the resected tumor confirmed it as a benign schwannoma (Figure 1c).

Discussion

Presacral schwannomas are rare accounting for 1-5% of spinal schwannomas [2-4]. Less than 30 cases of presacral schwannoma have been reported in English literature [5]. Klimo et al. [6] classified these tumors into 3 categories: Type 1 tumors are confined to the sacrum. Type 2 tumors originate within the sacrum but erode the wall of the sacrum and extend into the adjacent spaces. Type 3 tumors are located predominantly in the presacral or retro peritoneal area [6].

Most of these tumors remain asymptomatic and are detected incidentally during imaging studies done for nonspecific symptoms. Because of their slow growing nature and their location in the presacral fossa they can reach very large size while being asymptomatic and can even erode the sacrum. MRI is more specific than computed tomography (CT) as it can better delineate the tumor as well as its relationship with surrounding structures including intra spinous extension [7].

Complete surgical resection remains the treatment of choice. In general, posterior approach is preferable for type 1 tumors; anterior approach is preferable for type 3 tumors and combined anterior and posterior approach is preferable for type 2 tumors [8]. Anterior approach may be either trans peritoneal or extra peritoneal. Majority of the previous reports have reported trans peritoneal approach although extra peritoneal approach has also been reported by few authors [9,10]. The present patient had type 3 tumor and hence was resected via anterior trans peritoneal approach.




Figure 1A: MRI pelvis, axial section showing soft tissue mass arising in the presacral area with no obvious intra spinous extension.

Figure 1B: Gross picture of specimen of the presacral tumor resected.

Figure 1C: Spindle cell tumor with indented wavy nuclei revealing focal palisade.

Conclusion

Presacral schwannoma are rare nerve sheath tumors. MRI is the most preferred imaging modality. Complete surgical resection can be performed safely via anterior transabdominal approach.

References

1. Getachew MM, Whitman GJ, Chew FS (1994) Retroperitoneal schwannoma. *AJR Am J Roentgenol* 163(6): 1356.
2. Lin CM, Kao CC, Lin TC, Cha TL, Wu ST (2010) Giant presacral schwannoma mimicking malignancy in a man. *Acta Chir Belg* 110(3): 387-389.
3. Karabulut Z, Besim H, Hamamcı EO, Bostanoğlu S, Erverdi N, et al. (2002) Sacral schwannoma: case report. *Turk Neuro chir* 12: 247-505.
4. Domínguez J, Lobato RD, Ramos A, Rivas JJ, Gómez PA, et al. (1997) Giant intrasacral schwannomas: report of six cases. *Acta Neurochir (Wien)* 139(10): 954-960.
5. Konstantinidis K, Theodoropoulos GE, Sambalis G, Georgiou M, Vorias M, et al. (2005) Laparoscopic resection of presacral schwannomas. *Surg Laparosc Endosc Percutan Tech* 15(5): 302-304.
6. Klimo P, Rao G, Schmidt RH, Schmidt MH (2003) Nerve sheath tumors involving the sacrum. Case report and classification scheme. *Neurosurg Focus* 15(2): E12.
7. Popuri R, Davies AM (2002) MR imaging features of giant pre-sacral schwannomas: a report of four cases. *Eur Radiol* 12(9): 2365-2369.
8. Mazreku A, Karaj A, Avdia I, Bilali S (2010) The presentation and management of presacral tumors. *Acta Chir Iugosl* 57(2): 55-59.
9. Rousseau MA, Pascal-Mousselard H, Lazenec JY, Saillant G (2005) The mini-invasive anterior extra peritoneal approach to the pelvis. *Eur J Surg Oncol* 31(8): 924-926.
10. Senoglu M, Bulbuloglu E, Demirpolat G, Altun I, Ciralik H (2010) The anterior extraperitoneal approach to the rare presacral/retroperitoneal schwannoma. *Bratisl Lek Listy* 111(10): 558-561.